



# REPRODUCTIVE SYSTEM

**Based on GROUP-IV Examination syllabus -prepared by  
G.SRIRAM**

**NOTE: Dear kalam achievers kindly read at lest 4 to 5 times you can easily understand..**


---

## Reproduction In Animals

- There are two modes by which animals reproduce. These are:
  1. Sexual reproduction, and
  2. Asexual reproduction.
- There are many organisms which do not reproduce (mules, sterile worker bees, infertile human couples, etc.).

## Sexual Reproduction

- The reproductive parts in animals produce **gametes** that fuse to form a **zygote**. It is the zygote which develops into a new individual. This type of reproduction beginning from the fusion of male and female gametes is called **sexual reproduction**.
- The male reproductive organs include a pair of **testes (singular, testis)**, **two sperm ducts** and a **penis**. The testes produce the male gametes called **sperms**.
- The female reproductive organs are a **pair of ovaries, oviducts (fallopian tubes)** and the **uterus**. **Ovary** produces female gametes called **ova (eggs)**.




- In human beings, a single matured egg is released into the oviduct by one of the ovaries every month. **Uterus** is the part where development of the baby takes place. Like the sperm, an egg is also a single cell.
- The first step in the process of reproduction is the fusion of a sperm and an ovum. When sperms come in contact with an egg, one of the sperms may fuse with the egg. Such fusion of the egg and the sperm is called **fertilization**.

## Internal and External Fertilization


- During fertilization, the nuclei of the sperm and the egg fuse to form a single nucleus. This results in the formation of a fertilized egg or zygote.
- Fertilization which takes place inside the female body is called **internal fertilization**. Internal fertilization occurs in many animals including humans, cows, dogs and hens.
- During spring or rainy season, frogs and toads move to ponds and slow flowing streams. When the male and female come together in water, the female lays hundreds of eggs.
- Unlike hen's egg, frog's egg is not covered by a shell and it is comparatively very delicate. A layer of jelly holds the eggs together and provides protection to the eggs.
- As the eggs are laid, the male deposits sperms over them. Each sperm swims randomly in water with the help of its long tail. The sperms come in contact with the eggs. This results in fertilization.
- This type of fertilization in which the fusion of a male and a female gamete takes place outside the body of the female is called **external fertilization**. It is very common in **aquatic animals** such as fish, starfish, etc.

### Asexual Reproduction

- In each hydra, there may be one or more bulges. These bulges are the developing new individuals and they are called **buds**. In hydra, the new individuals develop as outgrowths from a single parent. This type of reproduction in which only a single parent is involved is called **asexual reproduction**. Since new individuals develop from the buds in hydra, this type of asexual reproduction is called **budding**.
- Another method of asexual reproduction is observed in the microscopic organism, amoeba. Reproduction in which an animal reproduces by dividing into two individuals is called **binary fission**. Apart from budding and binary fission, there are other methods by which a single parent reproduces the young ones.



**Fig. 9.11 : Budding in Hydra**



**Fig. 9.12** : Binary fission in Amoeba


### Human Reproductive System

- The reproductive events in humans include
  1. formation of gametes (**gametogenesis**), i.e., **sperms** in males and **ovum** in females,
  2. transfer of sperms into the female genital tract (**insemination**),
  3. fusion of male and female gametes (**fertilisation**) leading to formation of **zygote**.
  4. formation and development of **blastocyst** and its attachment to the uterine wall (**implantation**),
  5. embryonic development (**gestation**) and
  6. delivery of the baby (**parturition**).


### Male Reproductive System

- The male reproductive system is located in the pelvis region. It includes a pair of **testes** along with **accessory ducts**, **glands** and the external **genitalia**.
- The testes are situated outside the abdominal cavity within a pouch called **scrotum**. The scrotum helps in **maintaining the low temperature** of the testes (2–2.5<sup>0</sup>C lower than the normal internal body temperature) necessary for spermatogenesis.

- Each testis has about 250 testicular lobules. Each lobule contains one to three highly coiled **seminiferous tubules** in which **sperms are produced**.
- Each seminiferous tubule is lined on its inside by two types of cells called **male germ cells (spermatogonia)** and **Sertoli cells**.
- The male germ cells undergo **meiotic divisions** finally leading to sperm formation, while Sertoli cells provide **nutrition** to the germ cells.
- The regions outside the seminiferous tubules called interstitial spaces, contain small blood vessels and interstitial cells or **Leydig cells**. Leydig cells synthesise and secrete testicular hormones called **androgens** [a male sex hormone, such as **testosterone**. Androgens stimulates or controls the development and maintenance of male characteristics].



**Figure 3.1(b)** Diagrammatic view of male reproductive system (part of testis is open to show inner details)




**Figure 3.1(a)** Diagrammatic sectional view of male pelvis showing reproductive system

- The male sex accessory ducts include **rete testis, vasa efferentia, epididymis** and **vas deferens**.
- The seminiferous tubules of the testis open into the vasa efferentia through rete testis.
- The vasa efferentia leave the testis and open into epididymis. The epididymis leads to vas deferens that ascends to the abdomen and loops over the urinary bladder. It receives a duct from **seminal vesicle** [gland that secrete many of the components of semen] and opens into urethra as the ejaculatory duct. These ducts **store and transport** the sperms from the testis to the outside through urethra.
- The urethra originates from the urinary bladder and extends through the penis to its external opening called urethral meatus.
- The penis is the male external genitalia. It is made up of special tissue that helps in erection of the penis to facilitate insemination. The enlarged end of penis called the glans penis is covered by a loose fold of skin called foreskin.
- The male accessory glands include paired seminal vesicles, a prostate [releasing a fluid component of semen] and paired bulbourethral glands.
- Secretions of these glands constitute the **seminal plasma** which is **rich in fructose, calcium and certain enzymes**.
- The secretions of bulbourethral glands also helps in the **lubrication** of the penis.

### Female Reproductive System

- The female reproductive system consists of a pair of **ovaries** along with a pair of **oviducts, uterus, cervix, vagina** and the external **genitalia** located in pelvic region.


- These parts of the system along with a pair of the mammary glands are integrated structurally and functionally to support the processes of ovulation, fertilisation, pregnancy, birth and child care.
- **Ovaries** are the primary female sex organs [testis in males] that produce the female gamete (**ovum**) [sperm in males] and several **steroid hormones** (ovarian hormones).
- The ovaries are located one on each side of the lower abdomen. Each ovary is connected to the pelvic wall and uterus by ligaments.
- Each ovary is covered by a thin epithelium which encloses the ovarian stroma. The stroma is divided into two zones – a peripheral cortex and an inner medulla.
- The **oviducts (fallopian tubes)**, **uterus** and **vagina** constitute the female accessory ducts.



**Figure 3.3 (b)** Diagrammatic sectional view of the female reproductive system

- Each fallopian tube extends from the periphery of each ovary to the uterus, the part closer to the ovary is the funnel-shaped infundibulum.
- The edges of the infundibulum possess finger-like projections called fimbriae, which help in **collection of the ovum after ovulation**. The infundibulum leads to a wider part of the oviduct called ampulla.
- The last part of the oviduct, isthmus has a narrow lumen and it joins the uterus. The uterus is single and it is also called womb. The shape of the uterus is like an inverted pear.
- It is supported by ligaments attached to the pelvic wall. The uterus opens into vagina through a narrow cervix. The cavity of the cervix is called cervical canal which along with vagina forms the birth canal.

- The wall of the uterus has three layers of tissue. The external thin membranous perimetrium, middle thick layer of smooth muscle, myometrium and inner glandular layer called endometrium that lines the uterine cavity.
- The **endometrium** undergoes cyclical changes during **menstrual cycle** while the **myometrium** exhibits strong contraction during **delivery** of the baby.
- The female external genitalia include mons pubis, labia majora, labia minora, hymen and clitoris.
- Mons pubis is a cushion of fatty tissue covered by skin and pubic hair. The labia majora are fleshy folds of tissue, which extend down from the mons pubis and surround the vaginal opening.




**Figure 3.3 (a)** Diagrammatic sectional view of female pelvis showing reproductive system

- The labia minora are paired folds of tissue under the labia majora. The opening of the vagina is often covered partially by a membrane called hymen.
- The clitoris is a tiny finger-like structure which lies at the upper junction of the two labia minora above the urethral opening.
- The hymen is often torn during the first **coitus (intercourse)**. However, it can also be broken by a sudden fall or jolt, insertion of a vaginal tampon, active participation in some sports like horseback riding, cycling, etc.
- In some women the hymen persists even after coitus. In fact, the presence or absence of hymen is **not a reliable indicator of virginity** or sexual experience.

- A functional mammary gland is characteristic of all female mammals. The mammary glands are paired structures (breasts) that contain **glandular tissue** and variable amount of fat.
- The glandular tissue of each breast is divided into 15-20 mammary lobes containing clusters of cells called **alveoli**. The cells of alveoli **secrete milk**, which is stored in the cavities (lumens) of alveoli. The alveoli open into mammary tubules.
- The tubules of each lobe join to form a mammary duct. Several mammary ducts join to form a wider mammary ampulla which is connected to lactiferous duct through which milk is sucked out.

### **Gametogenesis**

- The primary sex organs – the **testis in the males** and the **ovaries in the females** produce gametes, i.e, **sperms** and **ovum**, respectively, by the process called **gametogenesis**.
- In testis, the immature **male germ cells (spermatogonia)** produce sperms by spermatogenesis that begins at puberty.
- The spermatogonia (sing. spermatogonium) present on the inside wall of seminiferous tubules multiply by **mitotic division** and increase in numbers. Each spermatogonium is **diploid** and contains 46 chromosomes.
- Some of the spermatogonia called primary spermatocytes periodically undergo **meiosis**.
- A primary spermatocyte completes the first meiotic division (reduction division) leading to formation of two equal, **haploid cells** called secondary spermatocytes, which have only 23 chromosomes each.
- The secondary spermatocytes undergo the **second meiotic division** to produce four equal, haploid spermatids.



**Figure 3.8** Schematic representation of (a) Spermatogenesis; (b) Oogenesis


What would be the number of chromosome in the spermatids? 23 chromosomes.

- The spermatids are transformed into spermatozoa (sperms) by the process called spermiogenesis. After spermiogenesis, sperm heads become embedded in the Sertoli cells, and are finally released from the seminiferous tubules by the process called spermiation.
- Spermatogenesis starts at the age of puberty due to significant increase in the secretion of **gonadotropin releasing hormone (GnRH)**. This, if you recall, is **ahypothalamic hormone**.
- The increased levels of GnRH then acts at the **anterior pituitary gland** and stimulates secretion of two **gonadotropins** – **luteinising hormone (LH)** and **follicle stimulating hormone (FSH)**.
- **LH** acts at the **Leydig cells** and stimulates **synthesis and secretion of androgens**. Androgens, in turn, stimulate the process of spermatogenesis.
- **FSH** acts on the **Sertoli cells** and stimulates secretion of some factors which help in the process of spermiogenesis.
- Sperm is a microscopic structure composed of a head, neck, a middle piece and a tail. A plasma membrane envelops the whole body of sperm.
- The sperm head contains an elongated haploid nucleus, the anterior portion of which is covered by a cap-like structure, acrosome. The acrosome is filled with enzymes that help fertilization of the ovum.

- The middle piece possesses numerous **mitochondria**, which produce energy for the movement of tail that facilitate sperm motility essential for fertilization.
- The human male ejaculates about 200 to 300 million sperms during a coitus of which, for normal fertility, at least 60 per cent sperms must have normal shape and size and at least 40 per cent of them must show vigorous motility.
- Sperms released from the seminiferous tubules, are transported by the accessory ducts.
- Secretions of epididymis, vas deferens, seminal vesicle and prostate are essential **maturation and motility of sperms**.
- The seminal plasma along with the sperms constitute the **semen**. The functions of male sex accessory ducts and glands are maintained by the **testicular hormones (androgens)**.
- The process of formation of a mature female gamete is called **oogenesis** which is markedly different from spermatogenesis.
- Oogenesis is initiated during the **embryonic development** stage when a couple of million gamete mother cells (oogonia) are formed within each fetal ovary; **no more oogonia are formed and added after birth**.
- These cells start division and enter into prophase-I of the meiotic division and get temporarily arrested at that stage, called **primary oocytes**.
- Each primary oocyte then gets surrounded by a layer of **granulosa cells** and is called the **primary follicle**.
- A large number of these follicles degenerate during the phase from **birth to puberty**. Therefore, at puberty only 60,000-80,000 primary follicles are left in each ovary.
- The primary follicles get surrounded by more layers of granulosa cells and a new theca and are called secondary follicles. The secondary follicle soon transforms into a tertiary follicle which is characterised by a fluid filled cavity called antrum.
- At this stage the primary oocyte within the **tertiary follicle** grows in size and completes its first **meiotic division**. It is an unequal division resulting in the formation of a large **haploid secondary oocyte** and a tiny first polar body.
- The secondary oocyte retains bulk of the nutrient rich cytoplasm of the primary oocyte.
- The tertiary follicle further changes into the mature follicle or **Graafian follicle**. The secondary oocyte forms a new membrane called zona pellucida surrounding it.
- The Graafian follicle now ruptures to release the secondary oocyte (**ovum**) from the ovary by the process called **ovulation**.

## Menstrual Cycle

- The reproductive cycle in the female primates (e.g. monkeys, apes and human beings) is called menstrual cycle. The first menstruation begins at puberty and is called **menarche**.
- In human females, menstruation is repeated at an average interval of about 28/29 days, and the cycle of events starting from one menstruation till the next one is called the **menstrual cycle**.




**Figure 3.9** Diagrammatic presentation of various events during a menstrual cycle

- One ovum is released (ovulation) during the **middle** of each menstrual cycle. The cycle starts with the menstrual phase, when menstrual flow occurs and it lasts for 3-5 days.
- The menstrual flow results due to breakdown of endometrial lining of the uterus and its blood vessels which forms liquid that comes out through vagina. Menstruation only occurs if the released ovum is **not fertilized**.
- Lack of menstruation may be **indicative of pregnancy**. However, it may also be caused due to some other underlying causes like stress, poor health etc.
- The menstrual phase is followed by the follicular phase. During this phase, the primary follicles in the ovary grow to become a fully mature Graafian follicle and simultaneously the endometrium of uterus regenerates through proliferation.

- These changes in the ovary and the uterus are induced by changes in the levels of pituitary and ovarian hormones.
- The secretion of gonadotropins (LH and FSH) increases gradually during the follicular phase, and stimulates follicular development as well as secretion of **estrogens** by the growing follicles.
- Both LH and FSH attain a peak level in the middle of cycle (about 14 th day). Rapid secretion of LH leading to its maximum level during the mid-cycle called LH surge induces rupture of Graafian follicle and thereby the **release of ovum** (ovulation).
- The ovulation (ovulatory phase) is followed by the luteal phase during which the remaining parts of the Graafian follicle transform as the corpus luteum.
- The corpus luteum secretes large amounts of **progesterone** which is essential for maintenance of the endometrium. Such an endometrium is necessary for implantation of the fertilised ovum and other events of pregnancy.
- During pregnancy all events of the menstrual cycle stop and there is no menstruation. In the absence of fertilisation, the corpus luteum degenerates. This causes disintegration of the endometrium leading to menstruation, marking a new cycle.
- In human beings, menstrual cycles ceases around 50 years of age; that is termed **asmenopause**.
- Cyclic menstruation is an indicator of normal reproductive phase and extends between menarche and menopause.


### **Fertilisation And Implantation**

- During copulation (coitus) semen is released by the penis into the vagina (insemination). The motile sperms swim rapidly, pass through the cervix, enter into the uterus and finally reach the ampullary region of the **fallopian tube**.
- The ovum released by the ovary is also transported to the ampullary region where fertilisation takes place.
- Fertilisation can only occur if the ovum and sperms are transported simultaneously to the **ampullary region**. This is the reason why not all copulations lead to fertilisation and pregnancy.



**Figure 3.10** Ovum surrounded by few sperms

- The process of fusion of a sperm with an ovum is called fertilisation. During fertilisation, a sperm comes in contact with the **zona pellucida** layer of the ovum and induces changes in the membrane that block the entry of additional sperms. Thus, it ensures that **only one sperm** can fertilise an ovum.
- The secretions of the acrosome help the sperm enter into the cytoplasm of the ovum through the zona pellucida and the plasma.



1 Transport of ovum, fertilisation and passage of growing embryo through fallopian tube

- Ovum surrounded by few sperm blastomeres is called a **morula**. The morula continues to divide and transforms into blastocyst as it moves further into the uterus.
- The blastomeres in the blastocyst are arranged into an outer layer called **trophoblast** and an inner group of cells attached to trophoblast called the inner cell mass. The trophoblast layer then gets attached to the **endometrium** and the inner cell mass gets differentiated as the **embryo**.
- After attachment, the uterine cells divide rapidly and covers the blastocyst. As a result, the blastocyst becomes embedded in the endometrium of the uterus. This is called **implantation** and it leads to pregnancy.


### In Vitro Fertilization

- Have you heard of test tube babies? In some women **oviducts are blocked**. These women are unable to bear babies because sperms cannot reach the egg for fertilization. In such cases, doctors collect freshly released egg and sperms and keep them together for a few hours for IVF or **In Vitro Fertilization (fertilization outside the body)**.
- In case fertilization occurs, the zygote is allowed to develop for about a week and then it is placed in the mother's uterus. Complete development takes place in the uterus and the baby is born like any other baby.
- Babies born through this technique are **called test-tube babies**. This term is actually misleading because babies cannot grow in test tubes.

### **Pregnancy And Embryonic Development**

- After implantation, finger-like projections appear on the trophoblast called chorionic villi which are surrounded by the uterine tissue and maternal blood.
- The **chorionic villi** and **uterine tissue** become interdigitated with each other and jointly form a structural and functional unit between developing embryo (foetus) and maternal body called **placenta**.
- The placenta facilitate the supply of oxygen and nutrients to the embryo and also removal of carbon dioxide and excretory/waste materials produced by the embryo.
- The placenta is connected to the embryo through an **umbilical cord** which helps in the transport of substances to and from the embryo.
- Placenta also acts as an **endocrine tissue** and produces several hormones like human chorionic gonadotropin (hCG), human placental lactogen (hPL), estrogens, progesterones, etc.
- In the later phase of pregnancy, a hormone called relaxin is also secreted by the ovary. Let us remember that hCG, hPL and relaxin are **produced in women only during pregnancy**.
- In addition, during pregnancy the levels of other hormones like estrogens, progesterones, cortisol, prolactin, thyroxine, etc., are increased several folds in the maternal blood.
- Increased production of these hormones is essential for supporting the fetal growth, metabolic changes in the mother and maintenance of pregnancy.
- Immediately after implantation, the inner cell mass (embryo) differentiates into an outer layer called ectoderm and an inner layer called endoderm. A mesoderm soon appears between the ectoderm and the endoderm [**triploblastic**]. These three layers give rise to all tissues (organs) in adults.
- It needs to be mentioned here that the inner cell mass contains certain cells called **stem cells** which have the potency to give rise to all the tissues and organs.
- The human pregnancy lasts 9 months. In human beings, after one month of pregnancy, the embryo's heart is formed. The first sign of growing foetus may be noticed by listening to the heart sound carefully through the stethoscope.
- By the end of the second month of pregnancy, the foetus develops limbs and digits. By the end of 12 weeks (first trimester), most of the major organ systems are formed, for example, the limbs and external genital organs are well developed.
- The first movements of the foetus and appearance of hair on the head are usually observed during the fifth month. By the end of about 24 weeks (end of second

trimester), the body is covered with fine hair, eye-lids separate, and eyelashes are formed. By the end of nine months of pregnancy, the foetus is fully developed and is ready for delivery.



**Figure 3.12** The human foetus within the uterus

### Parturition And Lactation


- The average duration of human pregnancy is about 9 months which is called the gestation period. Vigorous contraction of the uterus at the end of pregnancy causes expulsion/delivery of the foetus. This process of delivery of the foetus (childbirth) is called **parturition**.
- Parturition is induced by a complex neuroendocrine mechanism. The signals for parturition originate from the fully developed foetus and the placenta which induce mild uterine contractions called foetal ejection reflex. This triggers release of **oxytocin** from the maternal **pituitary**.
- Oxytocin acts on the uterine muscle and causes stronger uterine contractions, which in turn stimulates further secretion of oxytocin. The stimulatory reflex between the uterine contraction and oxytocin secretion continues resulting in stronger and stronger contractions. This leads to expulsion of the baby out of the uterus through the birth canal – parturition.
- Soon after the infant is delivered, the placenta is also expelled out of the uterus. The mammary glands of the female undergo differentiation during pregnancy and starts producing milk towards the end of pregnancy by the process called **lactation**. This helps the mother in feeding the newborn.

- The milk produced during the initial few days of lactation is called **colostrum** which contains several **antibodies** absolutely essential to develop resistance for the new-born babies.
- Breast-feeding during the initial period of infant growth is recommended by doctors for bringing up a healthy baby.

### Summary

- Humans are sexually reproducing and viviparous.
- The male reproductive system is composed of a pair of testes, the male sex accessory ducts and the accessory glands and external genitalia.
- Each testis has about 250 compartments called **testicular lobules**, and each lobule contains one to three highly coiled **seminiferous tubules**.
- Each seminiferous tubule is lined inside by spermatogonia and Sertoli cells.
- The spermatogonia undergo meiotic divisions leading to **sperm formation**, while Sertoli cells provide nutrition to the dividing germ cells.
- The **Leydig cells** outside the seminiferous tubules, synthesise and secrete testicular hormones called **androgens**.
- The male external genitalia is called penis.
- The female reproductive system consists of a pair of ovaries, a pair of oviducts, a uterus, a vagina, external genitalia, and a pair of mammary glands.
- The ovaries produce the female gamete (ovum) and some **steroid hormones** (ovarian hormones).
- Ovarian follicles in different stages of development are embedded in the stroma.
- The oviducts, uterus and vagina are female accessory ducts.
- The uterus has three layers namely perimetrium, myometrium and endometrium.
- The female external genitalia includes mons pubis, labia majora, labia minora, hymen and clitoris.
- The mammary glands are one of the female secondary sexual characteristics.
- Spermatogenesis results in the formation of sperms that are transported by the male sex accessory ducts.
- A normal human sperm is composed of a head, neck, a middle piece and tail.
- The process of formation of mature female gametes is called oogenesis.

- The reproductive cycle of female primates is called menstrual cycle.
- Menstrual cycle starts only after attaining sexual maturation (puberty).
- During ovulation only one ovum is released per menstrual cycle.
- The cyclical changes in the ovary and the uterus during menstrual cycle are induced by changes in the levels of **pituitary** and **ovarian hormones**.
- After coitus, sperms are transported to the junction of the isthmus and ampulla, where the sperm fertilizes the ovum leading to formation of a diploid zygote.
- The presence of X or Y chromosome in the sperm determines the sex of the embryo.



- The zygote undergoes repeated **mitotic division** to form a blastocyst, which is implanted in the uterus resulting in pregnancy.
- After nine months of pregnancy, the fully developed foetus is ready for delivery.
- The process of childbirth is called parturition which is induced by a complex neuroendocrine mechanism involving cortisol, estrogens and **oxytocin**.
- Mammary glands differentiate during pregnancy and secrete milk after child-birth.
- The new-born baby is fed milk by the mother (lactation) during the initial few months of growth.

Human seminal plasma, the fluid part of semen, is produced by contributions from the: i. Urethra ii. Prostate iii. Seminal vesicle iv. Bulbourethral gland

i and iii

i and iv

i, iii and iv

**ii, iii and iv**

### Question 2

Which one of the following statement is correct?

Spermatogenesis is the formation of sperm cells.

Spermatogenesis occurs in the seminiferous tubules.

Follicle-stimulating hormone (FSH) indirectly stimulates spermatogenesis.

**All of these are correct**

### Question 3

Which one of the following statements is incorrect about menopause?

Generally occurs between the ages of 45 and 55

Causes the cessation of the female reproductive cycle

**Anterior pituitary no longer produces FSH and LH**

Ovaries slowly reduce their secretion of estrogen

### Question 4

Which one of the following statement is correct about sertoli cells also called as nurse cells?

Found in adrenal cortex and secrete adrenaline

Found in ovaries and secrete progesterone

Found in pancreas and secrete cholecystokinin

**Found in Seminiferous tubules and provide nutrition of germ cells**

### Question 5

Which one of the following statements about morula in humans is correct?

It has more cytoplasm and more DNA than an uncleaved zygote

**It has almost equal quantity of cytoplasm as an uncleaved zygote but much more DNA**

It has far less cytoplasm as well as less DNA than in an uncleaved zygote

It has more or less equal quantity of cytoplasm and DNA as in uncleaved zygote

### Question 6

Foetal ejection reflex in human female is induced by

Differentiation of mammary glands

Pressure exerted by amniotic fluid

Release of oxytocin from pituitary

**Fully developed foetus and placenta**

### Question 7

Which one of the following is the most likely root cause why menstruation is not taking place in regularly cycling human female?

Retention of well-developed corpus luteum

Fertilisation of the ovum

Maintenance of the hypertrophical endometrial lining

**Maintenance of high concentration of sex-hormones in the blood stream**

### Question 8

The correct sequence of spermatogenetic stages leading to the formation of sperms in a mature human testis is

Spermatogonia-spermatid-spermatocyte-sperms

Spermatocyte-spermatogonia-spermatid-sperms

**Spermatogonia-spermatocyte-spermatid-sperms**

Spermatid-spermatocyte-spermatogonia-sperms

### Question 9

Correct sequence of embryo development:

**Gamete → Zygote → Morula → Blastula → Gastrula**

Gamete → Zygote → Blastula → Morula → Gastrula

Gamete → Neurula → Gastrula

Gamete → Neurula → Morula

### Question 10

Which one of the following statements is incorrect about menstruation?

The beginning of the cycle of menstruation is called menarche

During normal menstruation about 40 ml blood is lost

**The menstrual fluid can easily clot**

At menopause in the female, there is especially abrupt increase in gonadotropic hormones

### Question 11

The hemoglobin of a human foetus:

**Has a higher affinity for oxygen than that of an adult**

Has a lower affinity for oxygen than that of the adult

Its affinity for oxygen is the same as that of an adult

Has only two protein subunits instead of four

### Question 12

What happens during fertilisation in humans after many sperms reach close to the ovum?

	Cells of corona radiata trap all the sperms except one
	Only two sperms nearest the ovum penetrate zona pellucida
	<b>Secretions of acrosome helps one sperm enter cytoplasm of ovum through zona pellucida</b>
	All sperms except the one nearest to the ovum lose their tails


### Question 13

Column I contains terms and column II contains definitions. Match them correctly and choose the right answer.

	Column I		Column II
<b>A</b>	Parturition	<b>p</b>	Attachment of zygote to endometrium
<b>B</b>	Gestation	<b>q</b>	Release of egg from Graafian follicle
<b>C</b>	Ovulation	<b>r</b>	Delivery of baby from uterus
<b>D</b>	Implantation	<b>s</b>	Duration between pregnancy and birth
<b>E</b>	Conception	<b>t</b>	Formation of zygote by fusion of the egg and sperm
		<b>u</b>	Stoppage of ovulation and menstruation
	A = q, B = s, C = p, D = t, E = r		
	A = s, B = r, C = p, D = t, E = q		
	A = t, B = p, C = q, D = r, E = s		
	<b>A = r, B = s, C = q, D = p, E = t</b>		

**Question 14**


The figure given below depicts a diagrammatic sectional view of the female reproductive system of humans. Which one set of three parts out of I – VI have been correctly identified?



(I) Perimetrium, (II) Myometrium, (III) Fallopian tube
(II) Endometrium, (III) Infundibulum, (IV) Fimbriae
<b>(III) Infundibulum, (IV) Fimbriae, (V) Cervix</b>
(IV) Oviducal funnel, (V) Uterus, (VI) Cervix

**Question 15**

Given below is a diagrammatic sketch of a portion of human male reproductive system. Select the correct set of the names of the parts labelled A, B, C, D.



	A	B	C	D
(1)	Ureter	Seminal vesicle	Prostate	Bulbourethral gland
(2)	Ureter	Prostate	Seminal vesicle	Bulbourethral gland

(3)	Vas deferens	Seminal vesicle	Prostate	Bulbourethral gland
(4)	Vas deferens	Seminal vesicle	Bulbourethral gland	Prostate

	Option (1)
	Option (2)
	Option (3)
	Option (4)